

NORMAL FORMS OF MORSE-BOTT FUNCTIONS WITHOUT SADDLES ON COMPACT ORIENTED SURFACES

Bohdan Feshchenko

(Institute of Mathematics of NAS of Ukraine, Kyiv, Ukraine)

E-mail: fb@imath.kiev.ua

Let M be a smooth compact and oriented surface, and denote by P a real line \mathbb{R} or a circle S^1 . Denote by $\mathcal{F}^0(M, P)$ a class of Morse-Bott functions without saddles on M with the value in P . This class of functions naturally arises in the study of homotopy type of stabilizers of Morse-Bott functions on surfaces with respect to the action of the group of diffeomorphisms by pre-composition, see details in [1]. It is known that this class is non-empty if M is diffeomorphic to one of the following list: a cylinder $S^1 \times [0, 1]$, a disk D^2 , a sphere S^2 , a torus T^2 .

There are some trivial examples of functions from $\mathcal{F}^0(M, P)$ that are easy to write by hand:

Example 1. Let $f_0 : M_0 \rightarrow P$ be a smooth function from \mathcal{F}^0

- (1₀) $M_0 = S^1 \times [0, 1] = \{(z, s) \mid z \in \mathbb{C}, |z| = 1, 0 \leq s \leq 1\}$ is a unit cylinder, and $f_0 : S^1 \times [0, 1] \rightarrow \mathbb{R}$ is given by $f_0(z, s) = s$,
- (2₀) $M_0 = D^2 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}$ is a unit 2-disk, and $f_0 : D^2 \rightarrow \mathbb{R}$ is given by $f_0(x, y) = x^2 + y^2$,
- (3₀) $M_0 = S^2 = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}$ is a unit sphere, and $f_0 : S^2 \rightarrow \mathbb{R}$ is given by $f_0(x, y, z) = z$,
- (4₀) $M_0 = T^2 = \{(w, z) \in \mathbb{C}^2 \mid |z| = |w| = 1\}$ is a unit 2-torus, and $f_0 : T^2 \rightarrow S^1$ is given by $f_0(w, z) = z$.

Note that these functions do not have critical circles. We will call them **prime functions**.

Our main result is the following theorem, see [2].

Theorem 2. *A function $f \in \mathcal{F}^0(M, P)$ admits the following decomposition*

$$f = \varkappa \circ f_0 \circ h^{-1} \tag{1}$$

where $h : M_0 \rightarrow M$ is a diffeomorphism, $f_0 \in \mathcal{F}^0(M_0, P)$ is a prime function, and a smooth function $\varkappa : f_0(M_0) \rightarrow P$ which satisfies the following conditions:

- (A) \varkappa has the only finite number of non-degenerated critical points,
- (B) \varkappa does not have critical points at $f_0(\Sigma_{f_0})$ and $f_0(\partial M)$,

where Σ_{f_0} is the set of critical points of f_0 . A factorization (1) is not unique and depends on the choice of h . In particular, if f has no critical circles, then \varkappa is a diffeomorphism.

REFERENCES

- [1] Bohdan Feshchenko. Homotopy type of stabilizers of circle-valued functions with non-isolated singularities on surfaces. *arXiv:2305.08255*, 9p., 2023
- [2] Bohdan Feshchenko. Normal forms of functions with degenerate singularities on surfaces equipped with semi-free circle actions. *arXiv:2412.18944*, 18p., 2024.