

ON CARATHÉODORY THEOREM FOR ORLICZ-SOBOLEV CLASSES

Zarina Kovba

(Zhytomyr Ivan Franko State University)

E-mail: victoriazehrer@gmail.com

Evgeny Sevost'yanov

(Zhytomyr Ivan Franko State University; Institute of Applied Mathematics and Mechanics,
Slov'yans'k)

E-mail: esevostyanov2009@gmail.com

All definitions and notions used below may be found in [1]. The following notation is used: the set of prime ends corresponding to the domain D is denoted by E_D , and the completion of the domain D by its prime ends is denoted by \overline{D}_P . Below $C(f, \partial D)$ denotes the cluster set of f on ∂D . The following result holds.

Theorem 1. *Let $n - 1 < \alpha \leq n$, let D and D' be bounded domains in \mathbb{R}^n , $n \geq 3$, let $Q : D \rightarrow [0, \infty]$ be a Lebesgue measurable function and let $\varphi : [0, \infty) \rightarrow [0, \infty)$ be an increasing function. Let D be a regular domain and let $f : D \rightarrow D'$ be an open discrete mapping in $W_{\text{loc}}^{1, \varphi}(D)$, $f(D) = D'$. In addition, assume that $C(f, \partial D) \subset E_*$ for some closed (in the topology of \mathbb{R}^n) set $E_* \subset \overline{D}'$ and $f^{-1}(E_*) = E$ for some closed (in the topology of D) subset $E \subset D$. In addition, assume that: 1) the set E is nowhere dense in D and D is finitely connected on $E \cup \partial D$, i.e., for any $z_0 \in E \cup \partial D$ and any neighborhood \tilde{U} of z_0 there is a neighborhood $\tilde{V} \subset \tilde{U}$ of z_0 such that $(D \cap \tilde{V}) \setminus E$ consists of finite number of components, 2) for any $P \in E_D := \overline{D}_P \setminus D$ and for every neighborhood U of P in \overline{D}_P there is a neighborhood $V \subset U$ in \overline{D}_P of P such that $V \cap D$ is connected and $(V \cap D) \setminus E$ consists at most of m components, $1 \leq m < \infty$, 3) all components of the set $D' \setminus E_*$ have a strongly accessible boundary with respect to α -modulus, 4) the function φ satisfies the following Calderon condition*

$$\int_1^\infty \left(\frac{t}{\varphi(t)} \right)^{\frac{1}{n-2}} dt < \infty. \quad (1)$$

5) Assume that $K_{I, \alpha}(x, f) \leq Q(x)$ a.e. whenever $K_{I, \alpha}(x, f)$ is the inner dilatation of f of the order α and, in addition, the condition $\int_0^{\delta(b)} \frac{dt}{t^{\frac{n-1}{\alpha-1}} q_b'^{\frac{1}{\alpha-1}}(t)} = \infty$ holds for every point $b \in \partial D$ and some $\delta(b) > 0$, where $q_b'(t)$ denotes the integral average of the function Q' under the sphere $S(b, t)$, while $Q'(x) = \begin{cases} Q(x), & Q(x) \geq 1, \\ 1, & Q(x) < 1. \end{cases}$ Then f has a continuous extension $\bar{f} : \overline{D}_P \rightarrow \overline{D}'$, moreover, $\bar{f}(\overline{D}_P) = \overline{D}'$.

This result is published in [2].

REFERENCES

- [1] Martio O., Ryazanov V., Srebro U. and Yakubov E. *Moduli in Modern Mapping Theory*. Springer Science + Business Media, LLC : New York, 2009.
- [2] Kovba Z., Sevost'yanov E. *On Carathéodory prime ends extension for unclosed Orlicz-Sobolev classes*. <https://arxiv.org/abs/2604.15026>.

Acknowledgements. This talk is supported by the National Research Foundation of Ukraine (Project “Analogues of Carathéodory and Koebe-Bloch theorems for Orlicz-Sobolev classes”, Project number 2025.02/0010).